

Paper #60973

Variability in Orthodontic Treatment Planning Using the Invisalign Appliance

[S. BAUMRIND](#)¹, R.L. BOYD¹, M. DERAKHSHAN², S. REHAGE², and D. CHENIN¹, ¹ University of the Pacific School of Dentistry, Berkeley, CA, USA, ² Align Technology, Inc, Santa Clara, CA, USA

Objective: To investigate differences in the preferred solutions of experienced clinicians for the correction of anterior crowding, a common problem in the treatment of dental malocclusions. **Methods:** Other than extraction, three main strategies are available for resolving anterior crowding in orthodontic treatment. These include lateral expansion, incisor proclination, or inter-proximal reduction in tooth mass. A fourth possible strategy, distal displacement of upper posterior teeth, is advocated by some clinicians. To investigate the preferences of experienced clinicians among these three approaches when treating with the InvisAlign appliance, a panel of twelve clinical orthodontists was randomly chosen from among the 200 clinicians most experienced with InvisAlign treatment. Thirteen representative cases treated earlier using the appliance were identified and their pre-treatment records were duplicated electronically. The records for each case were re-evaluated independently by each of six blinded randomly chosen members of the panel, each of whom re-planned treatment for six cases. The matrix of cases by clinicians was designed to maximize the number of cases in which any two clinicians re-planned treatment for the same case. (All twelve clinicians evaluated the 13th case first for calibration and as a control.) **Results:** The table below summarizes the distribution of favored solutions for all cases. Two clinicians each considered the InvisAlign appliance to be inappropriate for the treatment of one case.

	Upper Arch			Lower Arch		
	Main Strategy	Secondary Strategy	Not Considered	Main Strategy	Secondary Strategy	Not Considered
Expansion	50	10	22	41	15	26
Proclination	14	43	25	16	45	21
Interproximal Reduction of tooth mass	19	35	28	32	38	11
"Distalization" of Upper Molars	9	----	75			

Conclusions: Almost all clinicians used each of the three solutions in some cases but the frequencies for different solutions differed markedly among clinicians and across cases.

Supported in part by Align Technology, Inc.

Abstract ID#: 60973

Password: 798231

Submitter's or presenter's Email: crilbaum@pacbell.net

Topic: Craniofacial Biology

Consider for Poster Presentation ONLY: N

The presenter is interested in becoming an oral session chair.

Keywords: Orthodontics, Human, Malocclusion, Treatment Planning

First Author

Presenting

Sheldon Baumrind
Orthodontics
University of the Pacific School of Dentistry
1525 Walnut Street
Berkeley, CA 94709
USA

Phone Number: 510-642-2709

Fax Number: 510-643-7829

Email: crilbaum@pacbell.net

Membership Number: P130825

Disclosure: This research project was supported in part by a grant from Align Technology, Inc. The presenter is a paid consultant for the company. However, the subject matter of this presentation is not related directly to the company's product

Second Author

Robert L. Boyd
Department of Orthodontics
University of the Pacific School of Dentistry
2155 Webster St
San Francisco, CA 94115
USA

Phone Number: 415-929-6690

Fax Number: 415-749-3390

Email: rboyd@sf.uop.com

Membership Number: P134455

Third Author

Mitra Derakhshan, DDS
Align Technology, Inc
881 Martin Avenue
Santa Clara, CA 95050
USA

Phone Number: 408-470-1333

Email: N/A

Fourth Author

Scott Rehage
Align Technology, Inc
881 Martin Avenue
Santa Clara, CA 95050
USA

Phone Number: 408-470-1439

Email: srebage@aligntech.com

Fifth Author

David Chenin, DDS

Orthodontics
University of the Pacific
2155 Webster Street
San Francisco, CA 94115
USA
Phone Number: 415-929-6555
Email: dchenin@pacific.edu